



## Case Report

# Early Prenatal Diagnosis of Congenital Ranula

Antonio Molero-Osorio<sup>1</sup>, Joel Santos-Bolívar<sup>1</sup>, Oscar Mencías<sup>2</sup>, Carlos Briceño-Pérez<sup>3</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, Maracaibo Ultrasound, <sup>2</sup>Department of Obstetrics and Gynecology, Materno Fetal Medicine and Obstetrics and Gynecology Services, Adolfo Pons Hospital, Social Security Venezuelan Institute, <sup>3</sup>Department of Obstetrics and Gynecology, Zulia University, Maracaibo, Venezuela



### \*Corresponding author:

Carlos Briceño-Pérez,  
Department of Obstetrics and  
Gynecology, Zulia University,  
Maracaibo, Venezuela.

cabripe@hotmail.com

Received : 14 October 19  
Accepted : 08 November 19  
Published : 13 December 19

DOI  
10.25259/AJS\_11\_2019

Quick Response Code:



## ABSTRACT

Congenital ranula (CR) is a rare tumor. Few cases of prenatal diagnosis (PD) of CR have been reported in the literature and diagnosis usually is beyond 20 weeks of gestation. We report a case of a pregnant patient with 16 weeks of gestation, where ultrasound (US) visualized in her fetus, a cystic tumor protruding in his oral cavity. Basis and complementary studies, including fetal echocardiography, were normal. Serial two-dimensional (2D) and three-dimensional (3D) US studies monitored fetal growth, amniotic fluid, progressive changes in the cystic mass, and maxillary movements. At 38.5 weeks of gestation, a cesarean section was performed. At 17<sup>th</sup> neonate days old, the neonate was done the simple and complete excision. Early PD of CR is possible. In this case was done as early as at 16 weeks of gestation. This is the first CR reported case with the earliest PD (16 week of gestation). We recommend that during the first trimester US, the oral cavity must be well explored, to rule out tumors.

**Keywords:** Congenital ranula, Mucocele, Congenital cystic malformation, Congenital high airway obstruction syndrome, Noninvasive prenatal diagnosis

## INTRODUCTION

Congenital ranula (CR) is a rare tumor, caused either by a mucus extravasation cyst secondary to obstruction and rupture of sublingual or minor salivary glands, or, less commonly, to a mucus retention cyst, originating from dilatation of an obstructed submandibular gland duct; and therefore surrounded by epithelium. Varies from 0.74 to 1.2/1000.<sup>[1-6]</sup> Few cases of prenatal diagnosis (PD) of CR have been reported in the literature and diagnosis usually is beyond 20 weeks of gestation.<sup>[1-3]</sup> We report a CR case, diagnosed as early as at 16 weeks of gestation, and review the literature.

## CASE REPORT

A 21-year-old pregnant patient, with no important personal or family history, G3 P2, during her routine prenatal control, at 16 weeks of gestation; by two-dimensional (2D) ultrasound (US) [Figure 1a], the open mouth was viewed in the fetus, and a rounded image of anechoic or cystic appearance, homogenous, and thin-walled, with no flow to the color Doppler, was visualized in the oral cavity. Three-dimensional (3D) US [Figure 1b] confirmed this image. There were no other congenital malformations or polyhydramnios. During prenatal care, she had a severe urinary infection that warranted hospitalization and parenteral treatment. She then continued antenatal care on an outpatient basis, her complementary studies including fetal echocardiography were normal and rejected the cytogenetic study. Serial 2D and 3D US monitored fetal growth, amniotic

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2019 Published by Scientific Scholar on behalf of American Journal of Sonography

fluid and progressive changes in the cystic mass, and observed maxillary movements and were unremarkable. US was performed using a GE Voluson 730 Expert and a Medison-Samsung Accuvix XG. At 38.5 weeks of gestation, an elective cesarean section was done; obtaining a male newborn, alive, 2.900 g, 49 cm, with Apgar scores of 8/10 and 9/10 at 1' and 5' of life. On clinical examination, the infant had a tumor on the ventral side of the tongue, 5 cm × 5 cm [Figure 2]. The patient was discharged 24 h postoperatively and the post-operative period was uneventful. A newborn was approached by a multidisciplinary team including obstetricians, neonatologists, and maxillofacial surgeons; he maintained the suction reflex and was fed orally without needing a nasogastric tube. At 17 days old, he had decreased 800 g and simple and



**Figure 1:** A 21-year-old woman at 16 weeks of gestation 2D ultrasound (top and bottom left), showed in the fetal oral cavity a rounded image of anechoic or cystic aspect, homogeneous, thinwalled, with no flow to the color Doppler assessment. 3D image (bottom right), showed the cystic mass in the fetal oral cavity.



**Figure 2:** Neonate's physical examination showed a cystic tumor on the ventral side of the tongue.

complete surgical excision was performed [Figure 3 top, middle and bottom left], the post-operative period was also uneventful [Figure 3 bottom center]. Four weeks later, the child was healthy, showed a normal-sized tongue, and had no feeding difficulties [Figure 3 bottom right]. Histological report confirmed the diagnosis.

## DISCUSSION

PD of congenital head and neck masses is becoming more frequent with the improvement of US technology, although it can be difficult to establish the etiology, which often is done by the histological examination.<sup>[1,7]</sup> CR may be diagnosed by prenatal US. With 2D US, is noted at the level of the floor of the mouth, a cystic, anechoic and avascular with color Doppler mass, which sometimes moves with fetal swallowing movements, displacing the tongue upward; that, in cases of large dimensions, can cause polyhydramnios.<sup>[3,5]</sup> The lesion lacks color Doppler flow inside and since it has elastic characteristics, it changes in shape with the reflex movements of suction. Other described PD method is 3D. US, which can play a role as adjuvant to 2D US, allowing a spatial evaluation of the relation between the mass and the oral cavity of the fetus. 3D US and magnetic resonance can also help diagnosis and differential diagnosis.<sup>[1]</sup> In the present case, diagnosis of the cystic lesion with 2D US was facilitated by the US findings of homogeneity, thin and regular walls, and an avascular mass to the color Doppler [Figures 1a and b]. The findings with 3D



**Figure 3:** At seventeen days of age, complete surgical excision of the cystic tumor was performed (top, middle and bottom left) in the neonate. Bottom center, the neonate on the first postoperative day. Bottom right, the neonate, 30 days after surgery.

US were similar to the previous ones. Correct PD is extremely important because, generally, such neonates present varying degrees of airway obstruction, which requires the presence of a multidisciplinary team at the time of birth.<sup>[1,5]</sup>

The literature reported that gestational age in which the PD is done is: Swiss Society of Neonatology,<sup>[8]</sup> 20 weeks; Onderoglu *et al.*<sup>[2]</sup> and Marques *et al.*,<sup>[5]</sup> 21 weeks; Pires *et al.*,<sup>[3]</sup> 25 weeks; Fernandez Moya,<sup>[4]</sup> 26 weeks; and Rios *et al.*,<sup>[1]</sup> 29 weeks. This case was diagnosed at 16 weeks of gestation.

Differential diagnosis must be done with epignathus, epulis, oropharyngeal teratoma, gingival cysts (palatine), hamartomas, lymphangiomas, thyroglossal duct cyst, congenital anomalies of the submandibular canal, heterotopic gastric cysts, and enterocystomas.<sup>[2-5]</sup> Epignathus is an unusual lesion estimated to affect between 1 in 35.000 and 1 in 200.000 live births. It is a benign teratoma mostly attached to the palatal or pharyngeal surface. Occasionally, these lesions may have an intracerebral extension. They are usually unresectable and affected infants usually die secondary to respiratory compromise. It is crucial that a timely tracheostomy is available during birth, while the fetomaternal circulation is uninterrupted.<sup>[2]</sup> The congenital epulis is a rare smooth mass, which occurs on the alveolar ridges, more common in the upper maxillary and in the female sex. In 10% of cases is a multiple lesion. It is a self-limiting lesion that responds to conservative excision.<sup>[2,9]</sup> The gingival or palatal cysts of the newborn are common and self-resolving. They originate from embryonic epithelial rests.<sup>[2]</sup> Vascular hamartomas represent proliferation of lymphatic or blood vessels.<sup>[2]</sup> Oral lymphangiomas can present superficially mostly on the tongue or they can less commonly present as deeper, poorly defined masses.<sup>[2]</sup>

Antenatal management involves US follow-up and early detection of the main obstetrics complications (polyhydramnios and preterm birth),<sup>[7]</sup> and other fetal morphological alterations. In our case, antenatal evolution was normal. US follow-up focused on fetal growth, amniotic fluid, maxillary movements and changes in the size, and position of cystic mass and showed no changes.

If possible, delivery should be at term, in a hospital at the tertiary level. The delivery route should be considered according to individual evolution, with a team of neonatologists and maxillofacial surgery. If the lesion is of large dimensions and/or obstruction of the upper airway is suspected, causing respiratory distress, before interrupting the fetoplacental connection, ex-uterus intrapartum treatment (EXIT) must be executed.<sup>[2,3,5]</sup> We discussed the mode of delivery and decided a cesarean section to avoid airways obstruction and rupture of the cyst. There was not necessary the aspiration or drainage of the cyst or the EXIT treatment.

Treatment of the ranula may be observation, aspiration, marsupialization, or surgical excision.<sup>[3,5]</sup> Cases of spontaneous

remission have been reported. The most appropriate and definitive treatment should be surgical excision, just as marsupialization or excision is chosen, with a recurrence risk of 5%.<sup>[3,5]</sup> The most common surgical complication is the rupture of the ranula during surgery, which does not increase the risk of recurrence.<sup>[3,5]</sup> In our case, simple surgical excision was performed, which evolved normally [Figure 3] and there was no rupture. It is thought that prenatal needle aspiration for tumor decompression prevents postnatal airway obstruction and obviates the EXIT procedure or neonatal intubation, especially if it is performed before birth.<sup>[3]</sup>

The most important neonatal complications are the loss of suction reflex, respiratory distress, language disorders, and interference with chewing and swallowing.<sup>[3,5]</sup> The present newborn decreased weight 800 g.

## CONCLUSIONS

We report the first CR case with the earliest PD (16 weeks of gestation) and recommend that during the first trimester US, oral cavity must be well explored, to rule out tumors.

## Acknowledgments

The authors acknowledge to Jonny Armas his valuable collaboration.

## Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

1. Rios LT, Júnior EA, Nardoza LM, Moron AF, Martins MG. Congenital ranula: Antenatal sonographic diagnosis. *Radiol Bras* 2012;45:300-1.
2. Onderoglu L, Saygan-Karamürsel B, Deren O, Bozdog G, Teksam O, Tekinalp G. Prenatal diagnosis of ranula at 21 weeks of gestation. *Ultrasound Obstet Gynecol* 2003;22:399-401.
3. Pires P, Pereira M, Machado L, Bonilla-Musoles F. Prenatal diagnosis of a ranula with 2- and 3-dimensional sonography and sonographically guided aspiration. *J Ultrasound Med* 2006;25:1499-502.
4. Fernandez Moya JM, Cifuentes Sulzberger S, Díaz Recaséns J, Ramos C, Sanz R, Perez Tejerizo G. Antenatal diagnosis and management of a ranula. *Ultrasound Obstet Gynecol* 1998;11:147-8.

5. Marques MI, Morais S, Coutinho S, de Castro O, Rei AI. Congenital ranula. *Acta Med Port* 2010;23:937-40.
6. Pontes FS, de Souza LL, Pedrinha VF, Pontes HA. Congenital ranula: A case report and literature review. *J Clin Pediatr Dent* 2018;42:454-7.
7. Rodríguez Tárrega E, Fuster Rojas S, Gómez Portero R, Roig Boronat S, Pérez Martínez G, Zamora Prado J, *et al.* Prenatal ultrasound diagnosis of a cyst of the oral cavity: An unusual case of thyroglossal duct cyst located on the tongue base. *Case Rep Obstet Gynecol* 2016;2016:7816306.
8. Berger TM. Swiss Society of Neonatology. Congenital Sublingual Cyst. Swiss Society of Neonatology. Available from: <https://pdfs.semanticscholar.org/f73d/fbf113a3bd7ff2672c2a2c69c8adcb6a6efa.pdf>. [Last accessed on 2019 Nov 27].
9. Braga-Tavares H, Santos H, M-Pinto I, Ramos M, de Sousa P. Congenital epulis. *Cir Pediatr* 2009;22:49-51.

**How to cite this article:** Molero-Osorio A, Santos-Bolívar J, Mencías O, Briceño-Pérez C. Early Prenatal Diagnosis of Congenital Ranula. *Am J Sonogr* 2019;2:7.